



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME VII

New names and the final members of new combinations are in **bold face type**

- Abena jamaicensis*, 244
Abies grandis, 121
Abronia umbellata, 64
Abrus precatorius, 340
Abutilon hirtum, 238; *periplocifolium*, 238
Acacia riparia, 178
Achyla, 307; *prolifera*, 307
Actaea alba, 64
Adiantum tenerum, 144
Aecidiella Triumfettae, 253
Aecidium, 66, 86, 170, 171, 253, 254; **abscedens**, 315; *Borreriae*, 315; *Cephalanthi*, 22; *Cestri*, 191; *circumscripturn*, 316; *Cissi*, 316; *Clematidis*, 74; *Clibadii*, 317; *compositarum Eupatorii*, 232; *decoloratum*, 317; *desmium*, 175; *detritum*, 254; *expansum*, 317; **favaceum**, 253; *Hyptidis*, 246-248; *Leonotidis*, 245; *luzoniense*, 254; *Nesaeae*, 86, 89; *passifloriicola*, 254; *Phyllanthi*, 254; *pulverulentum*, 315; *Randiae*, 315; *Rivinae*, 235; *roseum*, 250; *solaniphilum*, 255; *Sorbi*, 48; *Tournefortiae*, 254; *Traxini*, 22; *tubulosum*, 255; *tucumanense*, 246; *Uleanum*, 255; *verbenicola*, 22; *Wedeliae*, 318
Aeschynomene americana, 330; *sensitiva*, 330
Aesculus glabra, 65
Agaricus, 95, 97-101; *adirondackensis*, 257; *albo-umbilicatus*, 257; *alveolatus*, 291; *anisarius*, 276; *apertus*, 257; *arvensis*, 100, 102, 152, 226; *auratocephala*, 258; *bicolor*, 105; *brumalis*, 264; *caespitosus*, 281; *campestris*, 96, 100-102, 152; *candicans*, 257; *Candolleanus*, 120; *carnosior*, 259; *catinus*, 258; *cellaris*, 276; *clavipes*, 259; *columbanus*, 259; *compressipes*, 259; *comtulus*, 102; *concavus*, 259; *connexus*, 260; *cyathiformis*, 259-261; *dealbatus*, 260; *dicolor*, 261; *diminutivus*, 300; *ectypus*, 106; *ectypoides*, 262; *erubescens*, 262; *facifer*, 281; *flaccidus*, 262; *flavidellus*, 263; *fumosus*, 263; *gallinaceus*, 263; *Gerardianus*, 263; *giganteus*, 264; *gilvus*, 264; *Hoffmani*, 264; *illudens*, 281; *infundibuliformis*, 264, 265; *inversus*, 264; *lacrymabundus*, 116; *leptolomus*, 265; *maculosus*, 265; *marmoreus*, 273; *maximus*, 265; *metachrous*, 261; *mollis*, 268; *monadelphus*, 281; *nebularis*, 268; *nudus*, 105; *odorus*, 276; *pachylus*, 280; *panaeolus*, 106; *peltigerinus*, 267; *personatus*, 105; *phyllophilus*, 268; *pileolarius*, 268; *pinarius*, 268; *pinophilus*, 269; *pithyophilus*, 269; *placomycetes*, 152; *Poculum*, 259; *porphyrellus*, 269; *purus*, 118; *radiozonarius*, 270; *reticeps*, 290, 291; *reticulatus*, 291; *rivulosus*, 270; *Rodmani*, 212; *setisedus*, 270, 271; *Sienna*, 271; *sinopticus*, 271; *socialis*, 272; *sordidus*, 106; *splendens*, 272; *squamulosus*, 274; *subhirtus*, 273; *submarmoreus*, 273; *subzonalis*, 274; *tabescens*, 281; *truncicola*, 276; *trullisatus*, 275; *umbilicatus*, 257; *velutinus*, 116; *vilescens*, 276; *violaceolamelatus*, 120; *violaceus*, 105; *virens*, 276; *viridis*, 276
Agaricus reticeps Mont., Notes on, 290
Ageratum conyzoides, 250
Agropyron, 74, 75; *Smithii*, 73, 78
Aklema petiolaris, 236
Albizzia Lebbeck, 146
Alchornea latifolia, 328, 329
Aleuria badia, 90; *pustulata*, 92; *sylvestris*, 93; *vesiculosa*, 91
Allium reticulatum, 66
Alpinia antillarum, 339
Ambrosia, 77; *artemisiaefolia*, 77; *psilostachya*, 76, 77, 88; *trifida*, 77, 88
Amelanchier, 78, 84; *canadensis*, 78, 83, 84, 88, 89; *vulgaris*, 83, 89
Amsonia salicifolia, 84
Andropogon, 71, 72, 88; *brevifolius*, 339
Anemia hirsuta, 326
Anemone, 75; *cylindrica*, 73, 88
Apios tuberosa, 66
Apiosporium, 25
Apocynum cannabinum, 65
Aquilegia, 74, 75; *caerulea*, 73; *canadensis*, 83; *flavescens*, 73, 83

- Arabis, 75, 88
 Arachis hypogea, 322
 Argemone mexicana, 334
 Argomyces **insulanus**, 179; Vernoniae, 179, 180
 Armillaria, 95, 98; mellea, 102, 131, 132, 152, 281; mellea exannulata, 281
 Arnica, 77
 Aronia, 83; arbutifolia, 83
 Arthur, J. C., Cultures of Uredineae in 1912, 1913, and 1914, 61; Uredinales of Porto Rico based on collections by F. L. Stevens, 168, 227, 315
 Arthur, J. C., & Fromme, F. D., The taxonomic value of pore characters in the grass and sedge rusts, 28
 Arundinaria, 22
 Asclepias, 40, 41; curassavica, 240-242; speciosa, 40, 41; verticillata, 66
 Ascomycetes, Structural parallelism between the spore-forms in the, 21
 Ascospores of Endothia parasitica, The effect of continued desiccation on the expulsion of, 126
 Ascotricha, 289
 Ascrista monticola, 340
 Ascus-producing species, Penicillium avellaneum, a new, 284
 Aspergillus, 284
 Aspidium marginale, 176
 Aster, 70, 75, 76, 79, 81, 83; Drummondii, 79, 81, 83; ericoides, 72; multiflorus, 72; Novae-Angliae, 72; paniculatus, 65, 69, 70, 75, 76, 79, 81, 83, 84, 87, 88; Tripoli, 70; Tweedyi, 81, 83, 88
 Averswaldia, 340; palmicola, 340
 Axonophus compressus, 326
 Bagnisiella, 338
 Baptisia bracteata, 66
 Basanacantha, 315
 Bauhinia, 185; divaricata, 185; pauciflora, 184, 185
 Bidens andicola, 195; leucantha, 195, 196; pilosa, 195, 196; tereticaulis, 195
 Bixa orellana, 148, 327
 Bjerkandera adusta, 299
 Blechnum Brownei, 249
 Bletia patula, 320; purpurea, 320
 Bolbitis varicolor, 214
 Bolete from California, A new, 44
 Boletinus, 100; pictus, 300, 305
 Boletus **californicus**, 215; griseus, 166; ornatipes, 166
 Borreria levis, 249, 250; verticillata, 249, 250
 Botryosphaeria, 24
 Boudiera, 199
 Bouteloua, 31
 Brodiaea, 63; pauciflora, 85, 89
 Bromus, 74, 75
 Brown, H. P., A timber rot accompanying Hymenochaete rubiginosa (Schr.) Lév., 1
 Buetteneria lateralis, 323
 Caesalpinia, 184
 Cajan Cajan, 186
 Cajanus indicus, 186, 187
 California, A new bolete from, 44
 Calkins, William Wirt, amateur mycologist, 57
 Callirhoe digitata, 64, 65, 81, 82; involucrata, 65, 81, 82
 Calonectria graminicola, 25
 Canavalia ensiformis, 176
 Canna, 234, 235; coccinea, 234; glauca, 234; indica, 234; portoricensis, 234
 Cardiospermum grandiflorum, 236; Halicacabum, 236; microspermum, 236
 Carduus, 77, Flodmanii, 77, 78
 Carex, 31, 62, 69, 70, 76, 78, 81, 83, 86; arctata, 67, 87; crinita, 67, 87; extensa, 70; filiformis, 86, 89; Goodenowii, 76; intumescens, 75, 88; maritima, 62; pubescens, 66, 67, 87; retrorsa, 69-87; scoparia, 69, 70, 87; tenuis, 67, 87; tribuloides, 83, 88; vulpinoidea, 79, 81, 87
 Cassia chamaecrista, 66; quinquangulata, 321
 Cauloglossum, 100
 Cayaponia americana, 192, 335; racemosa, 192
 Cenchrus echinatus, 228, 229, 339; myosuroides, 339; viridis, 228, 229
 Centella asiatica, 334
 Cercospora, 41; clavata, 41
 Ceriomyces affinis, 300; auriporus, 300; communis, 44; crassus, 132, 151; ferruginatus, 152, 300, 306; miniato-olivaceus, 152; nebulosus, 300; retipes, 166; scaber, 152; subglabripes, 300; viscidus, 300, 305
 Cerotelium Canavaliae, 169, 176
 Cerrera unicolor, 299
 Cestrum laurifolium, 191; macrophyllum, 191
 Chaetochloa brevispica, 231; imberbis, 231; onurus, 231; purpurascens, 231; scandens, 228; setosa, 227, 231; verticillata, 230
 Chaetomium, 289
 Chamaecyparis thyoides, 83, 86, 89

- Chamaesyce hirta*, 190; *hypericifolia*, 190; *prostrata*, 190; *serpens*, 190
Chanterel aurantiacus, 152, 300;
Chantarellus, 152; *infundibuliformis*, 300, 305; *umbonatus*, 300, 305
 Characters in the grass and sedge rusts, The taxonomic value of pore, 28
Chelone glabra, 64, 65
Chlorophyllum Molybdites, 152
Chrysopsis villosus, 72
Cicuta maculata, 83
Cissus sicyoides, 147, 316
Citromyces, 134
Claudopus, 34-37; *depluens*, 36, 37, 290; **mephiticus**, 290; *nidulans*, 290, 300; **subdepluens**, 36, 37
Claudopus, A new mephitic, 290
Claudopus, A parasitic species of, 34
Clavaria, 305; *cinerea*, 299; *flava*, 151; *fusiformis*, 299; *muscigena*, 49; *pinophila*, 299; *pistillaris*, 299
Clematis, 75; *Douglasii*, 73, 83; *Drummondii*, 82, 83, 88; *ligusticifolia*, 73, 74, 88; *virginiana*, 73, 74, 83
Clibadium erosum, 317; *surinamense*, 317
Clitocybe, 256, 262, 266, 277, 280, 291, 301; *adirondackensis*, 257, 258; *aggregata*, 280; *albicastanea*, 278; *albidula*, 257; *albiformis*, 278; *albo-umbilicata*, 257; *ampla*, 263; *angustissima*, 280; *aperta*, 257; *aquatica*, 281; *Arnoldi*, 271; *atrialba*, 278; *avellanealba*, 278; *biformis*, 258, 265; *Broadwayi*, 280; *brumalis*, 261; *brunnescens*, 278; *caespitosa*, 258; *candicans*, 257; *candida*, 271, 280; *catina*, 258; *centralis*, 257; *cerrusata*, 280; *chrysocephala*, 258; *clavipes*, 259, 266, 278, 301; *columbana*, 259; *compressipes*, 259; *con-cava*, 259; *connexa*, 260; *cuticolor*, 278; *dealbata*, 258, 260, 266, 275, 278; *dealbata sudorifica*, 274; **di-color**, 260, 278; *difformis*, 280; *ditopoda*, 273, 280; *ditopus*, 259; **Earlei**, 261; *eccentrica*, 261, 279, 301; *ectypa*, 280; *ectypoides*, 262, 263; *elixa*, 280; *erubescens*, 262; *fellea*, 262; *flaccida*, 262; *flavidella*, 263; *fragrans*, 280; *fumosa*, 263; *fuscipes*, 263; *gallinacea*, 263; *geotropa*, 265, 280; *Gerardiana*, 263; *gigantea*, 264; *griseifolia*, 278; *Harperi*, 278; **hiemalis**, 264; *hirneola*, 280; *Hoffmani*, 264; *hondensis*, 278; *illudens*, 115, 131, 132, 152, 259, 281; *incrustedata*, 280; *infundibuliformis*, 257, 258, 264, 273, 301; *in-ornata*, 280; *inversa*, 258, 262, 264, 265, 272, 274, 278; *leptoloma*, 265; *maculosa*, 264, 265; *marginata*, 282; *maxima*, 264, 265; *media*, 266; *megalospora*, 157, 266; *metachroa*, 260, 261; *mexicana*, 280; *microspora*, 279; *monadelpha*, 281; *morbifera*, 260, 266, 275; *multiceps*, 152, 266, 267, 272; *multiformis*, 267; *murinifolia*, 278; *nebularis*, 34, 266, 268; *niveicolor*, 280; *nobilis*, 267; *obobata*, 280; *oculata*, 278; *odora*, 277; *opaca*, 280; *oreades*, 278; *oregonensis*, 278; *parasitica*, 281; *parilis*, 280; *Peckii*, 278; *peltigernia*, 267; *phyllophila*, 268; *phyllophiloides*, 214, 268; **pileo-laria**, 268; *pinariaria*, 268; *pinophila*, 269; *pithophylla*, 269; *porphyrella*, 269; *pruinosa*, 269, 280; *pulcherrima*, 270; *pusilla*, 279; *radi-ozonaria*, 270; *rancidula*, 270; *regularis*, 270; *revoluta*, 282; *rivulosa*, 270, 280; *robusta*, 271, 280; *rubro-tincta*, 280; *setiseda*, 271; *sinopica*, 269, 271, 272, 277, 279, 301; *sino-picoides*, 271; *socialis*, 272, 279; *sphaerospora*, 282; *splendens*, 272; *stipitata*, 278; *subcandicans*, 278; *subconcava*, 271, 272; **subconnexa**, 272; *subcyathiformis*, 273; *subdi-topoda*, 273, 301; *subfumosipes*, 278; *subhirta*, 273; *subinversa*, 278; *subinvoluta*, 280; **submar-morea**, 273; *subnigricans*, 274; *sub-similis*, 273; *subsocialis*, 279; **sub-squamata**, 272, 274; *sudorifica*, 260, 266, 274; *subzonalis*, 274; *sulphu-rea*, 275; **tenebricosa**, 275; *testa-ceoflava*, 280; *Trogii*, 260; *troyana*, 280; *trullaeformis*, 280; *trul-lisata*, 275; *truncicola*, 265, 276; *tuba*, 280; *tumulosa*, 280; *variabilis*, 278; *vilescens*, 276; *violaceifolia*, 278; *virens*, 260, 276, 277, 301; *washingtonensis*, 278
Clitocybe in North America, The genus, 256
Clitocybe megalospora, The validity of, 157
Clitoria cajanifolia, 189; *glycinoides*, 189; *rubiginosa*, 189
Clusia rosea, 336
Coccoloba, 145
Coccomyces, 25, 27; *hiemalis*, 45; *lutescens*, 45; *prunophorae*, 45
Coleosporium, 84, 169; *Elephantopodis*, 171; *Ipomoeae*, 80, 172; *Plumieriae*, 169, 172; *Solidaginis*, 80; *Vernoniae*, 80, 84, 89
 Collections by F. L. Stevens, Uredi-

- nales of Porto Rico based on, 168, 227, 315
- Colletotrichum*, 23, 38, 39, 40, 149; **destructivum**, 38; **salmonicolor**, 40; **solanicum**, 39; *Trifolii*, 38
- Colletotrichum* and *Phoma*, New species of, 38
- Collomia*, 77; *linearis*, 77, 88
- Collybia*, 291, 301; *acervata*, 301, 305; *confluens*, 301; *dryophila*, 301, 305; *longipes*, 132; *maculata*, 301, 306; *platyphylla*, 115, 118, 301; *radicata*, 157, 158, 301, 305; *retigera*, 291; *scabriuscula*, 301; *strictipes*, 301; *tuberosa*, 131, 132, 301, 302
- Colocasia*, 144
- Coltricia perennis*, 34, 299; *tomentosa*, 299
- Commelina elegans*, 182, 329; *nudiflora*, 183; *virginica*, 182, 329, 330
- Conard, Henry S., The structure and development of *Secotium agaricoides*, 94
- Continued desiccation on the expulsion of ascospores of *Endothia parasitica*, The effect of, 126
- Convolvulus*, 193; *nodiflorus*, 192
- Coprinus*, 98, 100; *atramentarius*, 34, 152; *comatus*, 152; *fimetarius*, 301; *micaceus*, 45, 102, 152, 301
- Cordia corymbosa*, 337
- Cordyceps militaris*, 298
- Coremium*, 134; *silvaticum*, 134
- Coriolus abietinus*, 299; *versicolor*, 299
- Corticium coeruleum*, 132
- Cortinarius*, 221, 223, 302; *anomalus*, 223; *armillatus*, 301; *erythrurus*, 223; 302; *lilacinus*, 302, 305; *purpurascens*, 302; **roseipallidus**, 221; *semisanguineus*, 302, 305
- Cortinellus*, 256; *decorus*, 164, 275
- Corydalis aurea*, 66
- Cosmos caudatus*, 195
- Cracca cinerea*, 177
- Crataegus cernuus*, 79, 89; *punctatus*, 78; *Pringlei*, 78, 88
- Craterellus*, 110; *borealis*, 110; *cornucopioides*, 299, 305
- Crepidotus*, 36, 37
- Crinipellis*, 156; *alnicola*, 156; *echinulata*, 156; *squamifolia*, 156; *sublivida*, 156; *zonata*, 270
- Cronartium Quercus*, 80, 89
- Crotalaria retusa*, 335
- Croton lucidus*, 147
- Crucibulum vulgare*, 305
- Cryptoporus*, 121, 123, 124; *volvatus*, 121, 123, 124, 125, 155
- Cryptoporus volvatus*, Notes on, 121
- Cryptosporella*, 24
- Cudonia lutea*, 298
- Cultures of Uredineae in 1912, 1913, and 1914, 61
- Cup-fungi, Photographs and descriptions of, —
I. *Peziza*, 90; II. *Sepultaria*, 197
- Cuphea*, 323; *Parsonsia*, 323
- Curcubitararia*, 27; *Laburni*, 24
- Cylindrosporium*, 25, 45
- Cyperus*, 231, 318; *cayennensis*, 231; *laevigatus*, 231; *mutisii*, 231; *polystachus*, 231; *radiatus*, 231; *sphacellatus*, 231; *surinamensis*, 231
- Cyphella*, 110
- Daedalea confragosa*, 299
- Daldinia concentrica*, 298
- Dasyscypha*, 9
- Dasyscypha foveolata*, 241
- Decodon*, 66; *verticillatus*, 86, 89
- Delphinium*, 73; *Geyeri*, 73
- Dendropanax arboreum*, 149
- Descriptions of cup-fungi, Photographs and, —
I. *Peziza*, 90; II. *Sepultaria*, 197
- Desiccation on the expulsion of ascospores of *Endothia parasitica*, The effect of continued, 126
- Desmodium axillare*, 188; *Scorpiurus*, 188
- Development of *Secotium agaricoides*, The structure and, 94
- Dicaeoma*, 28; *canaliculatum*, 231; *ciliata*, 319; *cubense*, 251; *deformatum*, 229; *Eleocharidis*, 232; *Paspali*, 230; *pulcherrimum*, 238; *radicans*, 319
- Dictyophora duplicata*, 153
- Dieffenbachia*, 144
- Dimerina*, 337; **Jacquiniæ**, 337
- Dimerium*, 335, 336, 337; **Cayaponiæ**, 335, 340; **grammodes**, 335, 337; **melioloides**, 336, 340; **Stevensii**, 337
- Dimerosporium*, 335; *lateritium*, 336
- Diodia rigida*, 250; *maritima*, 250
- Diorchidium leve*, 230
- Dioscorea*, 320; *grandiflora*, 320
- Dirca aecia*, 74; *palustris*, 64, 65
- Doellingeria umbellata*, 65
- Dolicholus reticulatus*, 186, 187; *texanus*, 186
- Dolichos*, 187; *Lablab*, 331; *reticulatus*, 187
- Dorstenia Contrajerva*, 327
- Dothidea grammodes*, 335; *perisporioides*, 335, 336; *seminata*, 335
- Dothidella grammodes*, 335
- Dryopteris patens*, 176, 325; *Poiteana*, 325

- Dulichium arundinaceum, 81, 87
 Durability of greenheart (*Nectandra Rodiaei* Schomb.), Tests on the, 204
- Edible and poisonous, Fungi, 151
 Effect of continued desiccation on the expulsion of ascospores of *Endothia parasitica*, The, 126
Elasmomyces, 99, 100; *mattirolianus*, 102
Eleagnus argentea, 65, 73
Eleocharis, 232; *cellulosa*, 232; *geniculata*, 232; *interstincta*, 232
Elephantopus angustifolius, 172; *carolinianus*, 84; *mollis*, 171, 172; *scaber*, 172
Eleutheranthera ruderalis, 251, 252
Elfvigia Brownii, 155, 215; *fomentaria*, 299; *megaloma*, 299
Elfvigiella, 110; *fasciata*, 110, 215
Elymus, 74, 75; *canadensis*, 73, 88; *virginicus*, 82, 88
Emilia sonchifolia, 251, 252
Endophyllum Rivinae, 235
Endoptychum, 99, 100
Endothia parasitica, 24, 27, 126, 130
Endothia parasitica, The effect of continued desiccation on the expulsion of ascospores of, 126
Entoloma, 105, 302; *cuspidatum*, 302; *sericeum*, 302; *strictius*, 302
Eocronartium typhuloides, 49
Eragrostis tephrosanthus, 180
Erigeron annuus, 66, 75
Eriosporangium Hyptidis, 247; *tucumanense*, 246
Erisyphc, 46, 328
Ernodea littoralis, 250
Erythrina glauca, 322, 323; *Lithospermae*, 146; *micropteryx*, 146, 323
Erythroxydon areolatum, 320, 321; *Coca*, 321; *havanense*, 320, 321
Eugenia buxifolia, 148; *Jambos*, 239
Eupatorium odoratum, 148; *polyodon*, 250; *villosum*, 251
Euphorbia corollata, 65; *cyparissias*, 65; *hirta*, 190; *petiolaris*, 236
Euthamia, 70; *graminifolia*, 70, 75, 83, 84, 87
Exogonium repandum, 149
 Expulsion of ascospores of *Endothia parasitica*, The effect of continued desiccation on the, 126
- Favolus variegatus*, 289
 Ferax group of the genus *Saprolegnia*, The, 307
Ficus, 174, 175; *Carica*, 174; *glom-erata*, 174; *laevigata*, 174
Fimbristylis, 232; *diphylla*, 232; *feruginea*, 232
 Fink, Bruce, William Wirt Calkins, amateur mycologist, 57
Fischeria crispiflora, 242
Fistulina, 155; *hepatica*, 151
 Fitzpatrick, Harry Morton, A parasitic species of *Claudopus*, 34
Flammula, 223; *penetrans*, 302; *polychroa*, 207, 208; *sapinea*, 302; *spumosa*, 302
Fomes Abramsianus, 215; *amarus*, 155; *annosus*, 132, 207, 208; **Brownii**, 215; *everhartii*, 207, 208; *fomentarius*, 207; *igniarius*, 155, 207, 208; *lobatus*, 207, 208; *nigricanus populinus*, 155; *pinicola*, 207, 208; *putearius*, 13; *robiniae*, 208; *roseus*, 207, 208, 299; *turbinatus*, 215; *ungulatus*, 299
Fomitiporia prunicola, 300
 Fromme, F. D., Arthur J. C., & The taxonomic value of pore characters in the grass and sedge rusts, 28
Fuirena, 319; *umbellata*, 319
Fuligo septica, 298
Fulvifomes, 110; *calcitratus*, 215; *Cedrelae*, 215; *cinchoensis*, 215; *dependens*, 215; *extensus*, 215; *grenadensis*, 215; *hydrophilus*, 215; *jamaicensis*, 215; *linteus*, 215; *melleicinctus*, 215; *pseudosenex*, 215; *sarcitus*, 215; *sublinteus*, 215; *suspectinatus*, 215. *Swieteniae*, 215; *troyanus*, 215; *Underwoodii*, 215; *yucatanensis*, 215
 Fungi edible and poisonous, 151
 Fungi, Illustrations of, — XX, 115; XXI, 163; XXII, 221
 Fungi, Luminescence in the, 131
 Fungi, Preliminary list of Upper St. Regis, 297
Fusarium conglutinans, 49
- Galera Hypnorum*, 302; *tener*, 302
 Garman, Philip, Some Porto Rican parasitic fungi, 333
 Genus *Clitocybe* in North America, The, 256
 Genus *Lepista*, The, 105
Gibberella, 27; *Saubineti*, 25
Gloeophyllum hirsutum, 300
Gloeoporus, 155
Gloeosporium, 23, 40
Glomerella, 23
Glycine reticulata, 186
Gnomonia, 24
Gossypium barbadense, 175; *brasiliense*, 175; *hirsutum*, 175
Gouania domingensis, 237, 329; *latifolia*, 238; *lupuloides*, 237, 329;

- polygama, 237, 329; tomentosa, 237, 329
- Grass and sedge rusts, The taxonomic value of pore characters in the, 28
- Greenheart (*Nectandra Rodiaei* Schomb.), Tests on the durability of, 204
- Grifola Berkeleyi, 46; frondosa, 151
- Grindelia squarrosa, 72
- Guarea trichiloides, 149
- Guettarda ovalifolia, 334
- Guignardia, 24
- Guilandina crista, 146
- Gutierrezia Sarothrae, 72, 88
- Gymnopilus **farinaceus**, 222; spumosus, 222
- Gymnosporangium, 48; Betheli, 78, 88; Botryapites, 83; clavariaeforme, 79, 88; durum, 78, 88; Ellisii, 86, 89; gracilens, 78, 88; Nelsoni, 78, 88; nidus-avis, 83, 89
- Gyrophila nimbata, 107
- Gyrophragmium, 100
- Hamaspora, 87
- Hapalopilus rutilans, 300
- Heald, F. D., & Studhalter, R. A., The effect of continued desiccation on the expulsion of the ascospores of *Endothia parasitica*, 126
- Hebeloma, 223; palustre, 214
- Helianthus angustifolius, 65
- Helicteres jamaicensis, 147
- Heliomyces, 156
- Helminthosporium, 109; gramineum, 49; teres, 49
- Helvella cochleata, 90, 91; Infula, 298
- Hemidiodia ocimifolia, 250, 315
- Herbarium, Marking types in the mycological, 108
- Herpotrichia, 210, 211; nigra, 23, 210, 211
- Herpotrichia nigra and associated species, Observations on, 210
- Heteropteryx, 240
- Heterotrichum cymosum, 337, 338
- Hibiscus militaris, 64, 82
- Humphrey, C. J., Tests on the durability of greenheart (*Nectandra Rodiaei* Schomb.), 204
- Hyalopsora, 170
- Hydnocystis, 197, 199
- Hydnum Caput-ursi, 151, 299; ochraceum, 299; repandum, 151; zonatum, 299
- Hydrophyllum capitatum, 73; Fendleri, 64, 73; virginicum, 64
- Hygrophorus, 259, 262; chlorophanus, 302; coccineus, 302, 305; miniatus, 302, 305; pratensis, 262
- Hymenaea, 321; Courbaril, 321
- Hymenochaete, 9; rubiginosa, 1-4, 13, 14-18
- Hymenochaete rubiginosa (Schrad.) Lév., A timber rot accompanying, 1
- Hymenoclea monogyra, 85, 89
- Hypholoma, 47, 98, 102; appendiculatum, 120, 152; Boughtoni, 117; Candolleum, 120; delineatum, 117; lacrymabundum, 116, 117; perplexum, 152; rigidipes, 117; rugocephalum, 117
- Hypomyces Ipomoeae, 25
- Hyptis atrorubens, 246; capitata, 247; latanifolia, 248; pectinata, 246; suaveolens, 246
- Illustrations of fungi—XX, 115; XXI, 163; XXII, 221
- Index to American Mycological Literature, 51, 113, 159, 217, 293, 341
- Indigofera Anil, 177; suffruticosa, 177
- Inga vera, 177
- Inocybe, 302; euthelella, 214; infida, 152
- Inonotus Leei, 155, 215; leprosus, 215; ludovicianus, 110; porrectus, 215; radiatus, 300
- Ipomoea, 80; acuminata, 243; Batatis, 172; carolina, 243; cathartica, 243; littoralis, 172; Nil, 172; pandurata, 80; stolonifer, 172; trichocarpa, 243; triloba, 243
- Iresine elatior, 235; paniculata, 235
- Iris versicolor, 64
- Ischnoderma fuliginosum, 300
- Jacquemontia nodiflora, 192, 193, 237; tamnifolia, 172
- Jacquinia barbasco, 337
- Jambos Jambos, 239
- Janipha Manihot, 190
- Jatropha Curcas, 331; gossypifolia, 331; Manihot, 190
- Juncus balticus, 76, 88
- Juniperus scopulorum, 78, 88; sibirica, 79, 89; utahensis, 78, 88; virginiana, 83, 89
- Klebahnna, 184, 196; **Bidentis**, 196
- Koeleria, 75; cristata, 75, 88
- Kuehneola, 169, 170, 174, 175; Fici, 174; Gossypii, 175
- Kyllinga caespitosa, 231; pumila, 231
- Laccaria, 256, 269; laccata, 275, 302, 305; striatula, 302
- Lachnea hemisphaerica, 299; scutellata, 299
- Laciniaria Langloisii, 84; punctata, 71, 84; scariosa, 77

- Lactaria, 302; camphorata, 302; deliciosa, 152; Gerardii, 302; Hibbardae, 225; lactiflua, 152, 166; ligulyota, 225, 302, 305; mucida, 302; oculata, 302; parva, 302; piperata, 152, 225, 302; plinthogala, 164; subdulcis, 302, 305; theiogala, 302; tominosa, 302, 306; turpis, 301, 302; varia, 302; **Volkertii**, 165
 Lactuca intybacea, 324
 Laetiporus speciosus, 132, 151
 Lamprospora, 199
 Lantana aculeata, 244; Camara, 243, 244, 334; crocea, 244; involucrata, 244; odorata, 244; stricta, 244; trifolia, 244
 Lasiacis divaricata, 180; Sloanei, 181; Swartziana, 181, 339;
 Lasiosphaeria Coulteri, 210
 Lentinellus, 156
 Lentinula, 156, 291; **reticeps**, 291
 Lentinus, 156; caespitosus, 281; cochleatus occidentalis, 216; lecomtei, 207, 208; lepideus, 119, 207; umbilicatus, 214
Lentodiellum, 157, 215, 216; **conca-vum**, 216
 Lentodium, 156; squamosum, 119
 Lenzites betulina, 207, 208; sepiaria, 207-209
 Leonotis nepetaefolia, 245
 Leotia lubrica, 299
 Lepargyrea canadensis, 65
 Lepiota, 119; americana, 152; amianthina, 118, 302; clypeolaria, 302, 305; fuscousquamea, 302; meleagris, 99, 102; naucina, 152, 302; procera, 152
 Lepista, 105; **domestica**, 105, 106; panaeola, 105, 106; personata, 105
 Lepista, The genus, 105
 Leptonia serrulata, 302
 Leptostroma, 23
 Limacella illinita, 302
 Lind's work on the Rostrup Herbarium, 42
 List of Upper St. Regis fungi, Preliminary, 297
 Lithospermum angustifolium, 71; officinale, 65
 Lophodermium, 23
 Luminescence in the fungi, 131
 Lycoperdon, 100; atropurpureum, 305; cyathiforme, 152, 305; gemmatum, 152, 305; pulcherrimum, 305; pyriforme, 305; separans, 305; subincarnatum, 305; Wrightii, 305
 Lycopus americanus, 71, 88
 Malvastrum, 82; coccineum, 66, 82
 Man, A new fungus, Phialophora verrucosa, pathogenic for, 200
 Manihot Manihot, 190; utilissima, 190
 Manisuris granularis, 230
 Marasmiellus, 156, 157; inconspicuus, 156; juniperinus, 156
 Marasmius, 155, 156, 301; **alnicola**, 156; campanulatus, 303; **distanti-folius**, 156; **echinulatus**, 156; **inconspicuus**, 156; **juniperinus**, 156; **niduliformis**, 156; oreades, 152; 303; rotula, 303; **squamifolius**, 156; **sublividus**, 156; **submulti-ceps**, 156
 Marasmius in North American Flora, Russula and, 155
 Marking types in the mycological herbarium, 108
 Medlar, E. M., A new fungus, Phialophora verrucosa, pathogenic for man, 200
 Meibomia, 189; adscendens, 335; axillaris, 188; Scorpiorus, 188; tortuosum, 188, 189
 Melampsoropsis, 175
 Melanoleuca, 118, 256, 267, 273; albissima, 274, 303, 306; fumidella, 165; melaleuca, 303; personata, 152; rutilans, 303; Russula, 222; sordida, 106
 Melanomma, 144
 Melanthera aspera, 194; canescens, 194, 195; deltoides, 195; hastata cubensis, 195
 Melasmia, 23
 Melothria guadalupensis, 192
 Menyanthes, 66
 Mephitic Claudopus, A new, 290
 Merulius, 155; lacrymans, 11, 206-209; tremellosus, 207, 208
 Mesosphaerum atrorubens, 246, 247; capitatum, 247; latanifolium, 248; pectinatum, 246, 247; spicatum, 246; suaveolens, 246, 247
 Metastelma lineare, 242; palustre, 243; parviflorum, 242; Schlechtendahlia, 242
 Miconia, 338; prasina, 337; Sintensii, 337, 338; laevigata, 333, 337
 Microcera, 23
 Mikania, 317; cordifolia, 317
 Milesia, 176; **columbiensis**, 175; **con-similis**, 176; **Kriegeriana**, 176
 Milesina columbiensis, 175; Kriegeriana, 176
 Mimosa ceratonia, 183, 184
 Mitracarpus portoricensis, 250
 Mitrula vitellina, 299
 Momisia iguanaea, 145
 Monadelphus, 256, 277, 281; caespi-

MacOwenites, 100

Macropodium macropus, 299

- tosus, 281; illudens, 258, 281, 282;
marginatus, 282; **revolutus**, 282;
sphaerosporus, 282
 Montagnites, 100
 Morchella esculenta, 151
 Muhlenbergia, 82, 181; Porteri, 81
 Murrill, W. A., Fungi edible and
 poisonous, 151; The genus Clitocybe
 in North America, 256; The gen-
 us Lepista, 105; Illustrations of
 Fungi—XX, 115, XXI, 163, XXII,
 221; Luminescence in the fungi,
 131; Marking types in the myco-
 logical herbarium, 108; A new bo-
 lete from California, 44; Prelimi-
 nary list of Upper St. Regis fungi,
 297
 Mutinus, 100
 Mycena, 280, 303; Leaiana, 303; pura,
 117, 260, 303
 Mycological herbarium, Marking types
 in the, 108
 Mycologist, amateur, William Wirt
 Calkins, 57
 Myrica cerifera, 87, 89
 Mytilidion, 210, 211; fusisporum, 211

 Napaea, 82; dioica, 82;
 Nectria discophora, 25; galligena, 25;
 megalospora, 336
 Neopectia Coulteri, 210, 211
 Nephrolepis rivularis, 175
 Neurocarpum cajanifolium, 189
 Neurolaena lobata, 252
 New ascus-producing species, Pencil-
 lium avellaneum, A, 284
 New Bolete from California, A, 44
 New fungus, Phialophora verrucosa,
 pathogenic for man, A, 200
 New mephitic Claudopus, A, 290
 News and Notes, 109, 212, 288
 and Reviews, 45, 155
 New species of Colletotrichum and
 Phoma, 38
 Nigredo, 28, 183, 192, 194; appendi-
 culata, 185; **columbiana**, 194; Com-
 melinae, 182; Dolicholi, 186; Erag-
 rostidis, 180; Hedysari-paniculati,
 188, 189; Helleriana, 192; leptoder-
 ma, 180; major, 181; Neuro-
 carpi, 189; proëminens, 190; Rhy-
 ncosporae, 182; Scleriae, 182
 North American Flora, Russula and
 Marasmius in, 155
 North America, The genus Clitocybe
 in, 256
 Notes on Agaricus reticeps Mont.,
 290
 Notes on Cryptoporus volvatus, 121
 Nyctalis, 34
 Observations on Herpotrichia nigra
 and associated species, 210
 Octospora pustulata, 92
 Oenothera grandiflora, 46; Tracyi, 46
 O'Gara, P. J., New species of Colleto-
 trichum and Phoma, 38
 Olyra latifolia, 229
 Omphalia 263, 304; campanella, 303;
 chrysophylla, 303; fibula, 303; um-
 bellifera, 304
 Onagra biennis, 75
 Onosmodium occidentale, 73
 Ophionectria, 27; coccicola, 23, 25
 Orton, C. R., Structural parallelism
 between spore-forms in the Asco-
 mycetes, 21
 Otidea, 91

 Panaeolus solidipes, 163
 Pandanus, 150
 Panellus, 156; flabellatus, 156; jala-
 pensis, 156; subcantharelloides, 156
 Panicum, 339; barbinode, 181; cap-
 illare, 65; divaricatum, 180; lana-
 tum, 181; maximum, 144; molle,
 181; ovalifolium, 229; Sloanei, 181;
 trichoides, 229; virgatum, 65
 Panus concavus, 216; **flabellatus**, 156;
 incandescens, 282; **jalapensis**, 156;
 meruliceps, 291; stypticus, 131,
 132; **subcantharelloides**, 156
 Parallelism between spore-forms in
 the Ascomycetes, Structural, 21
 Parasitic fungi, Some Porto Rican,
 333
 Parasitic fungi—I, Studies in Porto
 Rican, 143
 Parasitic species of Claudopus, A, 34
 Parodiella, 335, 337; melioloides, 336
 Paspalum, 230, 339; compressum,
 326; conjugatum, 319; fimbriatum,
 230; Helleri, 230, 326; Humboldtii-
 anum, 319, 326; orbiculatum, 230;
 paniculatum, 230, 326; pilosum,
 230; plicatulum, 230, 326; Schre-
 berianum, 230; Underwoodii, 339
 Passiflora rubra, 254
 Pathogenic for man, A new fungus,
 Phialophora verrucosa, 200
 Paxillus, 105; atrotomentosus, 304;
 involutus, 152, 304, 305; Lepista,
 105; panuoides, 304
 Peltigera, 267
 Pelvetia, 213
 Penicillium, 134, 135, 136, 138, 284;
 africanum, 135, 142; **avellaneum**,
 284; brevicaula, 134; candidum,
 284; claviforme, 134; duclauxi, 134,
 140, 141; expansum, 135; funicu-
 losum, 142; luteum, 135, 136, 137,
 140, 284; pinophilum, 135, 136, 140,

- 142; *purpurogenum*, 135, 136, 140, 142; *purpurogenum* var. *rubri-sclerotium*, 137; *rugulosum*, 140, 141, 142; *silvaticum*, 134
- Penicillium avellaneum*, a new ascus-producing species, 284
- Penicillium luteum-purpurogenum* group, The, 134
- Peridermium*, 79, 84; *carneum*, 80, 84; *Cerebrum*, 80, 214; *fusiforme*, 62, 79, 80, 84, 89;
- Petitita domingenis*, 333
- Peziza*, 90, 197; *abietina*, 299; *arenicola*, 197; *assimilata*, 92; *badia*, 90, 91, 93; *bufonia*, 93; *cochleata*, 90, 91; *heterophylla*, 73; *pustulata*, 92, 93; *sepulta*, 197, 198; *sylvestris*, 93; *tanacetifolia*, 64; *umbrina*, 92; *vesiculosa*, 91, 93
- Phaeolus sistotremaoides*, 300
- Phakopsora*, 173, 175; *Phyllanthi*, 332; *Vitis*, 173
- Phallus*, 100
- Phaseolus*, 186, 187; *adenanthus*, 185; *lunatus*, 335; *prostratus*, 187; *vulgaris*, 185
- Phialophora***, 202; *verrucosa*, 202, 203
- Phialophora verrucosa*, pathogenic for man, A new fungus, 200
- Philadelphus coronarius*, 78, 88; *Ke-teleerii*, 78, 88
- Philibertella clausa*, 243
- Philibertia*, 243
- Pholiota*, 304; *candicans*, 152; *Johnsoniana*, 226; *praecox*, 214; *squarrosa*, 304
- Phoma*, 41, 143; ***rostrata***, 41
- Phoma*, New species of *Colletotrichum* and, 38
- Photographs and descriptions of cup-fungi—
I. *Peziza*, 90; II. *Sepultaria*, 197
- Phragmidium*, 175
- Phragmites*, 22
- Phyllachora*, 337, 338; *Andropogonis*, 339, 340; *dalbergiicola* perforans, 340; *graminis*, 338, 339, 340; ***nitens***, 339; *paulensis*, 340; *perforans*, 340; *peribebuyensis*, 337, 338, 340; *Renalmiae*, 339; *sphaerosperma*, 339
- Phyllanthus distichus*, 332; *grandifolius*, 332; *nobilis*, 254
- Phyllosticta*, 143; ***adianticola***, 144; ***araliana***, 148; ***bixina***, 148; ***bo-ringuensis***, 147; ***cissicola***, 147; *Colocasiae*, 144; *colocasicola*, 144; ***commelinicola***, 144; *divergens*, 146; *Erythrinae*, 146; ***erythrini-***
cola, 146; ***Eugeniae***, 148; *eupatori-*
cola, 148; *glaucispora*, 149; ***guani-***
censis, 146; *Guareae*, 149; *Ipo-*
moeae, 149; *Lagenariae*, 149; ***mo-***
misiana, 145; *nericola*, 149; *Nerii*,
149; ***pandanicola***, 150; ***Panici***, 144;
Pithecolobii, 145; ***Pithecolobii mo-***
nensis, 145; ***portoricensis***, 147;
Sacchari, 144; *Sechii*, 149; ***Ste-***
vensii, 147
- Physalis pubescens*, 66
- Physopella*, 170, 173, 175, 330; *Aes-*
chynomenis, 330; *Fici*, 174; *ficina*,
174; *Vitis*, 173
- Picea*, 210; *Engelmanni*, 210
- Pieters, A. J., The *ferax* group of the
genus *Saprolegnia*, 307
- Pinus*, 169; *palustris*, 80, 89; *taeda*,
79, 80, 84, 89
- Piptoporus suberosus*, 300
- Pithecolobium Unguis-cati*, 145
- Pityrogramma calomelanos*, 325, 334
- Pleospora*, 25, 27, 49
- Pleuropus abortivus*, 152
- Pleurotopsis*, 156; *niduliformis*, 156
- Pleurotus*, 36, 37, 291; *caespitosus*,
281; ***cubensis***, 156; *lampas*, 282;
noctilucens, 282; *olearius*, 282; ***ori-***
zabensis, 156; *sapidus*, 152; ***sub-***
barbatulus, 156
- Plicaria badia*, 90; *pustulata*, 92
- Plowrightia*, 25
- Pluchea odorata*, 324; *purpurascens*,
324, 325
- Plumieria alba*, 173; *Krugii*, 172; *ob-*
tusa, 172; *rubra*, 173
- Pluteus alveolatus*, 291; *cervinus*, 152,
304
- Poinsettia heterophylla*, 190
- Poisonous, Fungi edible and, 151
- Polymarasmius*, 156, 157; *submulti-*
ceps, 156
- Polyplocium*, 100
- Polyporus*, 34, 35; *adustus*, 207; *arcu-*
larius, 214; *caudicinus*, 132; *ele-*
grans, 300; ***graminicola***, 215; *Leei*,
215; *Marbleae*, 215; *McMurphyi*,
155; *obtusus*, 207, 208; *perennis*,
34-37; ***porrectus***, 215; *resinosus*,
207, 208; *Schweinitzii*, 214; *sul-*
phureus, 207, 208; *volvatus*, 125;
Zelleri, 155
- Polystictus hirsutus*, 207, 208, 209;
prolificans, 208; *versicolor*, 207,
208
- Pore characters in the grass and
sedge rusts, The taxonomic value
of, 28
- Porodaedalea Pini*, 111, 132, 300
- Porothelium*, 155
- Porto Rican parasitic fungi, Some,
333

- Porto Rican parasitic fungi—I, Studies in, 143
- Porto Rico based on collections by F. L. Stevens, Uredinales of, 168, 227, 315
- Preliminary list of Upper St. Regis Fungi, 297
- Prospodium appendiculatum, 178
- Psalliotia, 101
- Pseudopeziza, 24
- Pseudotsuga taxifolia, 121
- Psidium, 240; Guajava, 239
- Psychotria patens, 323
- Pteris, 334
- Puccinia, 22, 28, 62, 67, 76, 170, 245, 322, 323; Agropyri, 73, 74, 82, 88; albiperidia, 67, 68, 69, 78, 87; alternans, 74; Andropogonis, 72; angustata, 64, 70, 71, 88; angustatoides, 62; appendiculata, 178; Arechavelatae, 236; Becki, 179; Blechi, 249; Boutelouae, 30; **Cameliae**, 227; Campulosi, 29; canaliculata, 231, 318; Cannae, 233; Caricis-Asteris, 32, 69, 76, 87; Caricis-Solidaginis, 69, 79, 87; Caricis-Strictae, 32; Cenchri, 32, 228; Chaetochloae, 230; compacta, 240, 241; **concrescens**, 240, 241; Convolvuli, 192; crassipes, 243; Cryptandri, 64; Cynanchi, 242; Cyperi, 231; deformata, 229; Dolichi, 186; Dulichii, 66, 84, 81, 87; Eleocharidis, 232; Eleutherantherae, 251; Ellisiana, 71, 72, 88; emaculata, 65; Emiliae, 251; epiphylla, 29, 32; esclavensis, 32; Euphorbiae, 236; extensicola, 70, 76, 79, 80, 81, 87; Fimbristylidis, 232; Fuirenae, 319; Gonolobi, 243; Gouaniae, 237, 329; gouaniicola, 238; graminis, 62; gregaria, 241; Grossulariae, 66, 67, 69, 76, 87; hemisphaerica, 324; heterospora, 238; Huberi, 229; Hyptidis, 247, 248; **insititia**, 248; Ipomoeae, 243; inflata, 236; Jambosae, 239, 240; karelica, 32; Latanae, 243; lateritia, 249; leonotidicola, 245; Leonotidis, 245; leptospora, 29; levis, 230; macropoda, 235; McClatchieana, 64; medellinensis, 246, 247, 249; Melicae, 30; minutissima, 86, 89; monoica, 75, 88; Muhlenbergiae, 81, 82, 88; Nesaeae, 86; nodosa, 85, 89; obliqua, 242, 243; oblitterata, 74; Pammelii, 65; Panici, 65; panicicola, 180; paradoxica, 29; Paspali, 230; peridermiospora, 22; polyculiformis, 32; Psidii, 239, 240; quadriporula, 76, 87; Raunkiaerii, 235; Ribesii-Caricis, 69; Rivinae, 235; Rompellii, 240; rosea, 250; rubigo-vera, 62; salviicola, 249; sanguinolenta, 240; **sclericola**, 232; Seymouriana, 22, 30; solida, 251, 252; sphaerospora, 242, 243; splendens, 85, 89; Sporoboli, 31, 32; Stipae, 72, 88; substriata, 230; Synedrellae, 251; tagetica, 251; Thaliae, 233; Thwaitessii, 238; tomipara, 74; tosta, 64, 66, 81, 82, 84, 88; triarculata, 30; Tridacis, 251; **tucumanensis**, 247; uniporula, 67, 68, 69, 78; Urbaniana, 244; Urticae, 32; Vernoniae, 179; Vernoniae-mollis, 179; versicolor, 30; vexans, 31, 32; Vilfae, 22; vulpinoides, 62, 63, 66, 79, 80, 84, 87; xanthopoda, 233
- Pucciniastrum, 169, 175
- Puccinosira, 253; pallidula, 170, 253;
- Pustularia vesiculosa, 91
- Pyropolyporus Abramsianus, 155, 215; cinnabarinus, 300; conchatus, 300; ignarius, 111, 300; ignarius nigricans, 111
- Quamoclit coccinea, 172
- Quercus Phellos, 80, 89; rubra, 79, 80, 89
- Rajania, 320; cordata, 320
- Randia, 315; aculeata, 315
- Ravenelia, 178, 184, 322, 329; Caesalpiniae, 183; capituliformis, 328; caulicola, 176; Indigoferae, 177; Ingae, 177; **Stevensii**, 178
- Resupinatus, 156; cubensis, 156; orizabensis, 156; subbarbatulus, 156
- Reticularia Lycoperdon, 298
- Rheosporangium aphanidermatum, 288
- Rhizina inflata, 213
- Rhizoctonia, 39, 48; solani, 48
- Rhodopaxillus, 105; panaeolus, 107; personatus, 105; parvidius, 106
- Rhynchospora, 62; aurea, 182; cyperoides, 182; distans, 182; micrantha, 182; polyphylla, 182
- Rhyncosia, 187; longeracemosa, 187; reticulata, 186; Senna, 187; texana, 186
- Rhytisma, 23
- Ribes, 69, 76; Cynosbati, 66, 78, 87; Rivina humilis, 235; octandra, 235
- Romell, Lars, Lind's work on the Rostrop Herbarium, 42
- Rosellinia melioloides, 336
- Rostkovites **californicus**, 44, 215; granulatus, 152, 300, 305; subaureus, 300, 305

- Rostrup Herbarium, Lind's work on the, 42
 Rostrupia Scleriae, 233
 Rot accompanying Hymenochaete rubiginosa (Schrad.) Lév., A timber, 1
 Ruellia strepens, 65
 Russula, 155, 156, 166, 221, 222, 225, 273, 304; albida, 304; bifida, 224; compacta, 304, 305; crustosa, 224; depallens, 304; emetica, 152, 304, 306; foetens, 152, 304, 306; furcata, 224; lutea, 304, 305; Mariae, 152; pectinatoides, 167; virescens, 152, 224
 Russula and Marasmius in North American Flora, 155
 Rusts, The taxonomic value of pore characters in the grass and sedge, 28
 Rytillix granularis, 230
 Sabicea aspera, 324
 Saccharum, 144
 Salvia occidentalis, 249
 Saprolegnia, 307; anisospora, 308; declina, 308, 312; dioica, 308; ferax, 307-314; hypogyna, 308; mixta, 307-314; monoica, 307-309, 312-314; Thureti, 307, 308; torulosa, 307, 308
 Saprolegnia, The ferax group of the genus, 307
 Sarcosphaera, 199
 Schizophyllum commune, 208
 Schizophyllum, 156
 Schlegelia brachyantha, 339
 Schroeteriaster, 170
 Scirpus atrovirens, 70, 88; Eriophorum, 71; fluviatilis, 83, 88; microcarpus, 64, 71; rubrotinctus, 64, 71; sylvaticus, 64, 71
 Scleria, 182, 233; Baldwinia, 233; hirtella, 233; pauciflora, 233; pterota, 182; setacea, 233; verticillata, 233
 Scleroderma, 167; aurantium, 153, 305; tenerum, 167
 Sclerotinia, 24
 Scopulariopsis, 134
 Scutiger griseus, 300; hispidellus, 155
 Scytinotus, 156; distantifolius, 156
 Seaver, Fred J., Observations on Herpotrichia nigra and associated species, 210; Photographs and descriptions of cup-fungi—I. Peziza, 90; II. Sepultaria, 197
 Sechium edule, 149
 Secotium, 98, 99-103; agaricoides, 94, 97, 99-102; coarotatum, 100; erythrocephalum, 99; Gueinzii, 100; Mattirolianus, 99; melanosporum, 100; olbium, 100, 103; Warnei, 94
 Secotium agaricoides, The structure and development of, 94
 Sedge rusts, The taxonomic value of pore characters in the grass and, 28
 Senecio Douglasii, 72; lugens, 72; obovatus, 81; spartioides, 72, 88
 Septoria, 333; aquilina, 334; asiatica, 334; Chelidonii, 334; Guettardae, 334, 340; Lantanae, 334, 340; Miconiae, 333, 340; Petitiae, 333, 340; Pityrogrammae, 334; Verbenae, 334
 Sepultaria, 197, 198, 199; arenicola, 198; Longii, 199
 Setaria scandens, 228; setosa, 227; verticillata, 230
 Sida, 239; glutinosa, 238; humilis, 238, 239; nervosa, 238; procumbens, 238; spinosa, 238; urens, 238, 239
 Sidalcea, 82
 Silphium perfoliatum, 77
 Sitodrepa panicea, 124
 Sium cicutaeifolium, 83, 88
 Solanum torvum, 255; tuberosum, 39
 Solenia, 155
 Solidago, 63, 75, 76, 79, 81, 83; canadensis, 65, 70, 72, 75, 79, 81, 83, 84, 87, 88; glaberrima, 79, 87; missouriensis, 79; mollis, 72, 79, 87; rugosa, 79, 83, 87
 Some Porto Rican parasitic fungi, 333
 Sparganophorum Vaillantii, 325
 Spartina, 22; Michauxiana, 77, 88
 Spathularia velutipes, 299
 Species of Claudopus, A parasitic, 34
 Species of Colletotrichum and Phoma, New, 38
 Spermocoe aspera, 250; tenuior, 250
 Sphaeralcea, 82; incana, 64, 81, 88; lobata, 64, 65, 82, 88
 Sphaeria canaliculata, 231; meliolioides, 336; perisporioides, 335, 336
 Spongipellis borealis, 300; fragilis, 110
 Spore-forms in the Ascomycetes, Structural parallelism between, 21
 Sporobolus, 82, 181; asperifolius, 81, 88; cryptandrus, 64; diandrus, 181; indicus, 181; longifolius, 66
 Stachytarpheta jamaicensis, 244; strigosa, 244
 Steironema ciliatum, 77; lanceolatum, 77
 Stemmodontia trilobata, 318
 Stenolobium Stans, 178
 Stereum, 2; fasciatum, 207, 208, 209; frustulosum, 13-17; gausapatum,

- 207, 208; *hirsutum*, 16; *rameale*, 207, 208
- Stigmaphyllon lingulatum*, 236; *periplocifolium*, 236; *Sagraeanum*, 236
- Stipa comata*, 72, 88
- Strobilomyces strobilaceus*, 152
- Stropharia*, 95, 98; *ambigua*, 103; *bi-lamellata*, 226; *coprinophila*, 34; *semiglobata*, 304
- Struchium Sparganophorum*, 325
- Structural parallelism between spore-forms in the Ascomycetes, 21
- Structure and development of *Secotium agaricoides*, The, 94
- Strumella*, 46; *coryneoides*, 46
- Studhalter, R. A., Heald, F. D., & The effect of continued desiccation on the expulsion of the ascospores of *Endothia parasitica*, 126
- Studies in Porto Rican parasitic fungi—I, 143
- Suillelus luridus*, 152, 300, 306
- Synedrella nodiflora*, 251
- Tagetes patula*, 251
- Taxonomic value of pore characters in the grass and sedge rusts, The, 28
- Tecoma Stans*, 178
- Tectella*, 156
- Teramnus uncinatus*, 331
- Tests on the durability of greenheart (*Nectandra Rodiaei* Schomb.), 204
- Tetrazygia*, 337, 338
- Thalia*, 234, 235; *geniculata*, 234
- Thalictrum*, 75; *alpinum*, 73; *dioicum*, 74, 83; *Fendleri*, 64, 73, 83
- Thelephora laciniata*, 299; *perdix*, 13
- Thom, Charles, The *Penicillium luteum-purpureogenum* group, 134
- Thom, Charles, & Turesson, G. W., *Penicillium avellaneum*, a new ascus-producing species, 284
- Timber rot accompanying *Hymenochaete rubiginosa* (Schr.) Lév., 1
- Tournefortia hirsutissima*, 254
- Tradescantia multiflora*, 183
- Trametes Pini*, 13; *abietis*, 13, 14, 17; *Pini*, 214; *robiniofila*, 207
- Tremella lutescens*, 299; *mycetophila*, 299
- Tremellodon gelatinosum*, 299
- Tricholoma*, 105, 256, 291; *cellare*, 277; *multipunctum*, 165; *nudum*, 105; *panaeolum*, 105, 107; *perso-natum*, 105, 270; *rancidulum*, 270; *Russula*, 222; *Sienna*, 277; *sordidum*, 601
- Tridax procumbens*, 252
- Trifolium carolinianum*, 85, 89; *hybrum*, 38; *pratense*, 38, 39
- Triopteris lingulata*, 236
- Trisetum*, 75; *subspicatum*, 75, 88
- Triumfetta*, 253; *grandiflora*, 253; *Lappula*, 253; *rhomboidea*, 253; *semitriloba*, 147, 253
- Tsuga heterophylla*, 121
- Turesson, G. W., Thom, Charles, & *Penicillium avellaneum*, a new ascus-producing species, 284
- Tylopilus felleus*, 151, 300, 306
- Types in the mycological herbarium, Marking, 108
- Typhula muscicola*, 49
- Tyromyces caesius*, 300; *chioneus*, 300; *graminicola*, 215; *guttulatus*, 300; *lacteus*, 300
- Upper St. Regis fungi, Preliminary list of, 297
- Urechites lutea*, 149
- Uredinales of Porto Rico based on collections by F. L. Stevens, 168, 227, 315
- Uredineae in 1912, 1913, and 1914, Cultures of, 61
- Uredo, 170, 330; *Aeschynomenis*, 330; *Agerati*, 250; *amaniensis*, 195; *Arachidis*, 322; *balaensis*, 249; *basipora*, 245; *bidenticola*, 195; *Bidentis*, 195; *biocellata*, 324; ***Bixae***, 327; *Cabreriana*, 322; *Cajani*, 187; *Cameliae*, 227, 228; *cancerina*, 246; *Cannae*, 233; *capituliformis*, 328; *Cephalanthi*, 324; *Cestri*, 191; *Chaetochloae*, 230; *Commelinaceae*, 182; *Commelinae*, 182; *Commelyneae*, 329; ***concors***, 330; *Cupheae*, 323; *Desmodii-tortuosi*, 189; *Dichromenae*, 319; *Dioscoreae*, 320; *Dolichi*, 186; *Erythroxylonis*, 320; *Eugeniarum*, 239, 240; ***fallaciosa***, 323; ***fenestrala***, 170, 332; *Fici*, 174; *ficicola*, 174; *ficina*, 174; *flavidula*, 239; *Fuirenae*, 319; *gemmata*, 193, 237; *Gossypii*, 175; *Gouaniae*, 329; *Gymnogonimmes*, 170, 325; *Hymenaeae*, 321; *ignobilis*, 181; *Jacquemontiae*, 193; *Janiphae*, 190; ***jatrophicola***, 331; *Kyllingiae*, 231; *leonotica*, 245, 246; *Leonotidis*, 245; ***lutea***, 321; *moricola*, 174; *Myrtearum*, 239, 240; *neurophylla*, 240; *nigropunctata*, 320; *nootkatensis*, 48; *pamparum*, 187; *paspalicola*, 319, 326; *Phyllanthi*, 332; ***proximella***, 324; *rubescens*, 327; ***sabiceicola***, 323; *Sparganophori*, 325; ***Stevensiana***, 326; *striolata*, 235; *subneurophylla*,

- 240; *superficialis*, 326; *Teramni*, 331; *vicina*, 325; *Violae*, 173; *Vitis*, 173
- Uromyces*, 22, 28, 31, 32, 170, 184, 240, 322, 323; *acuminatus*, 77, 88; *appendiculatus*, 185, 187; *Arachidis*, 322; *Bidentis*, 195, 196; ***Caesalpiniae***, 183; *Cestri*, 191; *columbianus*, 194; *Commelinae*, 182, 330; ***densus***, 196; *dichrous*, 190; *Dolicholi*, 186, 187; *elegans*, 85, 89; *Eragrostidis*, 180; *Euphorbiae*, 189; *euphorbiicola*, 190; *gemmatus*, 192, 237; *Hedysari-paniculati*, 188; *Hellerianus*, 192; ***ignobilis***, 181; *insularis*, 189; *jamaicensis*, 184; ***Janiphae***, 190; *Jatrophae*, 191; *Junci*, 76, 88; *leptodermus*, 180; *major*, 181; *malvacearum*, 238; *malvicola*, 238; *Neurocarpi*, 189; *pedatatus*, 72; *perigynius*, 75, 76, 83, 88; *pianhyensis*, 227, 325; *pictus*, 238; *Poae*, 32; *proëminens*, 189; *pulcherrimus*, 238; *quadripodula*, 76; *Rhyncosporae*, 65, 182; *Scirpi*, 83, 88; *Scleriae*, 182, 233; *Sidae*, 238; *solidus*, 188; *Sporoboli*, 66; *Thwaitesii*, 238; *uniporulus*, 32, 67, 68
- Vaginata agglutinata*, 152; *plumbea*, 152, 304, 305; *plumbea strangulata*, 305
- Vagnera stellata*, 66
- Valerianodes jamaicensis*, 244; *strigosa*, 244
- Validity of *Clitocybe megalospora*, The, 157
- Value of pore characters in the grass and sedge rusts, The taxonomic, 28
- Venenarius Caesareus*, 152; *cothurnatus*, 152; *phalloides*, 152, 305, 306; *Frostianus*, 305; *muscarius*, 152, 305, 306; *rubens*, 152; *solitarius*, 152; *spretus*, 152
- Venturia*, 25, 26, 27
- Vermicularia*, 39
- Vernonia*, 80, 84; *albicaulis*, 179; *boringuensis*, 180; *fasciculata*, 80, 84, 89; *longifolia*, 179; *phyllostachya*, 180
- Vigna*, 186, 187; *repens*, 185; *vexillata*, 185
- Vincetoxicum*, 243
- Viola*, 71, 72; *cucullata*, 71, 88; *Nuttallii*, 71, 88; *premulaefolia*, 71; *striata*, 72
- Vitis*, 173; *vinifera*, 173
- Volvaria Loveiana*, 34
- Wallrothiella Arceuthobii*, 289
- Wedelia*, 227, 325; *carnosa*, 318; *lanceolata*, 325; *reticulata*, 227; *trilobata*, 318
- Wissadula periplocifolia*, 238, 239
- Work on the Rostrup Herbarium, Lind's 42
- Xanthium*, 318; *canadense*, 77
- Xylaria Hypoxylon*, 131, 132; *polymorpha*, 299
- Xylopiia*, 241
- Young, Esther, Studies in Porto Rican parasitic fungi, 143
- Zanthoxylum americanum*, 66
- Zeller, Sanford M., Notes on *Cryptoporus volvatus*, 121